

SUPPLEMENTAL LABELING

Sulfomet™ Extra Herbicide

EPA Reg. No. 352-622-85588

FOR WEED CONTROL IN HYBRID POPLAR PLANTATIONS IN THE STATE OF NEW MEXICO

DIRECTIONS FOR USE

Sulfomet™ Extra herbicide controls specific weeds to aid in the establishment and growth of hybrid poplar plantings.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

SULFOMET™ EXTRA HERBICIDE USE RATES AND RESTRICTIONS

Site Preparation — Application Before Transplanting For hybrid poplar, apply 1 to 3 ounces per acre of Sulfomet™ Extra herbicide. Use 2 to 3 ounces per acre of Sulfomet™ Extra herbicide for heavy weed infestations and where maximum residual control is desired. Use 1 to 2 ounces per acre of Sulfomet™ Extra herbicide for light weed infestations or when small diameter cuttings have been planted. Allow a minimum of 3 days between application and planting. Limit the first use to a small area to determine the selectivity of Sulfomet™ Extra herbicide on specific clones. Sulfomet™ Extra herbicide must be activated by rainfall or overhead irrigation before weeds become well established. Use of Sulfomet™ Extra herbicide may cause temporary chlorosis (yellowing) or a small reduction in tree height during the year of use.

Release — Application After Transplanting

For hybrid poplar, apply 1 to 3 ounces per acre of Sulfomet™ Extra herbicide. Use 2 to 3 ounces per acre of Sulfomet™ Extra herbicide for heavy weed infestations and where maximum residual control is desired. Use 1 to 2 ounces per acre of Sulfomet™ Extra herbicide for light weed infestations or when small diameter cuttings have been planted.

Note: Apply only to trees which have been established for a minimum of 1 year. Apply when the trees are dormant and avoid contact of the spray with green buds or tissue as injury to the trees may result. Avoid applications during the period when the hybrid poplar are actively growing; from bud-swell in the spring to leaf drop in the fall. Limit the first use to a small area to determine the selectivity of Sulfomet™ Extra herbicide on specific clones.

Sulfomet™ Extra herbicide must be activated by rainfall or overhead irrigation before weeds become well established. Use of Sulfomet™ Extra herbicide may cause temporary chlorosis (yellowing) or a small reduction in tree height during the year of use.

Sold by Agsurf Corporation

© 2008 Agsurf Corporation, 1209 Orange Street, Wilmington, Delaware, 19801

Product information: www.agsurf.com

All rights reserved.

TANK MIXES

Sulfomet™ Extra herbicide can be tank mixed with other products that are registered for use on hybrid poplars and where the labeled method of application and timing of application are the same as for Sulfomet™ Extra herbicide.

Specific Weed Problems

Kochia and Russian Thistle

Since biotypes of kochia and Russian thistle are known to be resistant to Sulfomet™ Extra herbicide, tank mixture combinations with herbicides having different modes of action should be used.

To slow the development of resistant biotypes, minimize kochia or Russian thistle forming mature seed.

IMPORTANT PRECAUTIONS

- Applications of Sulfomet™ Extra herbicide made to hybrid poplar trees that are suffering from loss of vigor caused by insects, diseases, drought, winter damage, animal damage, excessive soil moisture, planting shock, previous agricultural practices, or other stresses, may injure or kill the trees.
- Applications of Sulfomet™ Extra herbicide made for release (trees present) should only be made after adequate rainfall has closed the planting slit and settled the soil around the roots following transplanting.
- If a surfactant is used with Sulfomet™ Extra herbicide, allowing the spray to contact tree foliage may injure or kill trees. The user assumes all responsibility for tree injury if a surfactant is used with Sulfomet™ Extra herbicide treatments applied after planting.
- Sulfomet™ Extra herbicide applications may result in damage and mortality to other species of trees when they are present on sites.

IMPORTANT

BEFORE USING THIS HERBICIDE, READ AND CAREFULLY NOTE THE CAUTIONARY STATEMENTS AND OTHER PROCEDURAL INFORMATION APPEARING ON THE EPA REGISTERED LABEL OR ON OTHER SUPPLEMENTAL LABELS.

This bulletin contains new or supplemental instructions for use of these products in combination which does not appear on the package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using THIS product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal product label.

R-845 090108 01-24-07

Sold by Agsurf Corporation

© 2008 Agsurf Corporation, 1209 Orange Street, Wilmington, Delaware, 19801

Product information: www.agsurf.com

All rights reserved.