

SUPPLEMENTAL LABELING

Chlormet™ Herbicide

EPA Reg. No. 352-445-85588

RECROPPING INTERVALS FOR GRASSES ON CONSERVATION RESERVE PROGRAM (CRP)

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

ROTATION INTERVAL FOR PLANTING GRASSES ON CONSERVATION RESERVE PROGRAM (CRP) ACRES

Whenever Chlormet™ herbicide has previously been used in wheat, barley, or fallow, the following grasses may be planted after the intervals specified in the tables below. **The planting of grass and legume mixtures is not recommended as injury to the legume may occur.**

- Bentgrasses
- Blue grama
- Bluestems — Big, Little, Plains, Sand, WW Spar
- Buffalograss
- Galleta
- Green needlegrass
- Green sprangletop
- Indiangrass
- Indian ricegrass
- Lovegrasses — Sand, Weeping
- Orchardgrass
(excluding Piaute)
- Prairie sandreed
- Sand dropseed
- Sheep fescue
- Sideoats grama
- Switchgrass
- Wheatgrasses — Crested, Intermediate, Pubescent,
Slender, Streambank, Tall, Thickspike, Western
- Wild-ryegrasses — Beardless, Russian

Sold by Agsurf Corporation

© 2008 Agsurf Corporation, 1209 Orange Street, Wilmington, Delaware, 19801

Product information: www.agsurf.com

All rights reserved.

ROTATION INTERVALS IN:

MT, ND, SD, and Northern WY:

<u>Soil pH*</u> <u>Grasses</u>	<u>Use Rate</u> <u>(oz/acre)</u>	<u>Minimum Interval for Planting</u>
7.5 or lower	2/10 - 3/10	4 months (all grasses)
7.6 to 7.9	2/10 - 3/10	4 months (Wheatgrasses only)

CO, NM, Southern WY:

<u>Soil pH*</u> <u>Grasses</u>	<u>Use Rate</u> <u>(oz/acre)</u>	<u>Minimum Interval for Planting</u>
7.9 or lower	2/10 - 3/10	2 months (all grasses)

NE, KS, OK, TX:

<u>Soil pH*</u> <u>Grasses</u>	<u>Use Rate</u> <u>(oz/acre)</u>	<u>Minimum Interval for Planting</u>
7.9 or lower	2/10 - 4/10	2 months (all grasses)
7.9 or lower	5/10	4 months (all grasses)

ID,OR, UT, WA:

<u>Soil pH*</u> <u>Grasses</u>	<u>Use Rate</u> <u>(oz/acre)</u>	<u>Minimum Interval for Planting</u>
7.9 or lower	2/10 - 4/10	2 months (all grasses)

BIOASSAY

A field bioassay must be completed before rotating to any crop not listed (See the Rotation Intervals table), or if the soil pH is not in the specified range, or if the use rate applied is not specified in the table, or if the minimum cumulative precipitation has not occurred since application.

Field Bioassay

To conduct a field bioassay, grow test strips of the crop or crops you plan to grow the following year in fields previously treated with Chlormet™ herbicide. Crop response to the bioassay will indicate whether or not to rotate to the crop(s) grown in the test strips. If a field bioassay is planned, check with your local Agricultural dealer or Agsurf representative for information detailing the field bioassay procedure.

Sold by Agsurf Corporation

© 2008 Agsurf Corporation, 1209 Orange Street, Wilmington, Delaware, 19801

Product information: www.agsurf.com

All rights reserved.

**IMPORTANT
BEFORE USING CHLORMET™ HERBICIDE, READ AND FOLLOW ALL
APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE
EPA-REGISTERED LABEL.**

This bulletin contains new or supplemental instructions for use of this product which do not appear on the EPA-registered package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

R-849 090108 07-22-98

Sold by Agsurf Corporation

© 2008 Agsurf Corporation, 1209 Orange Street, Wilmington, Delaware, 19801

Product information: www.agsurf.com

All rights reserved.